

Variability in life-history switch points across and within populations explained by Adaptive Dynamics

Pietro Landi^{1,*}

James R. Vonesh^{2,3}

Cang Hui^{1,4}

1. Theoretical Ecology Group, Department of Mathematical Sciences, Stellenbosch University, Matieland 7602, South Africa;

2. Center for Environmental Studies, Virginia Commonwealth University, Richmond, Virginia 23284-2012;

3. Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Matieland 7602, South Africa;

4. Mathematical and Physical Biosciences, African Institute for Mathematical Sciences, Muizenberg 7945, South Africa;

* Corresponding author; e-mail: landi@sun.ac.za

Journal of the Royal Society Interface

Supplementary figure




Figure S1: Scenarios for time of maturation under different juvenile initial sizes computed with logistic growth function $w(\tau) = g(w_0, \tau) = w_0 w_K \exp(r\tau) / (w_K + w_0(\exp(r\tau) - 1))$. Orange interval: internal Evolutionary Stable Strategy (ESS) $0 < \bar{\tau} < 1$. Parameters as in figure 4a in the main text except $w_K = 10$. Dark green interval: branching point (within-population variability). Light green interval: either branching point or boundary strategy $\bar{\tau} = 0$ (across- and within-population variability). Red interval: direct development. Solid line: convergence stable equilibria (attracting strategies). Dashed line: convergence unstable equilibria (repelling strategies). Thick line: convergence stable but evolutionary unstable equilibria (branching points). TC: transcritical bifurcation. BR: branching bifurcation ($B = 0$ in equation (6) in the main text), turning selection disruptive. SN: saddle-node bifurcation.