

## A-Amylase immobilization on amidoximated acrylic microfibers activated by cyanuric chloride

Yaaser Q. Almulaiky<sup>1,2</sup>, Faisal M. Aqlan<sup>3</sup>, Musab Aldhahri<sup>4,5</sup>, Mohammed Baeshen<sup>6</sup> Tariq Jamal Khan<sup>7</sup>, Khalid A. Khan<sup>8</sup>, Mohamed Afifi<sup>6,9</sup>, Ammar AL-Farga<sup>1</sup>, Mohiuddin Khan Warsi<sup>1</sup>, Mohammed Alkaled<sup>6</sup>, Aisha A.M. Alayafi<sup>6</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Science, University of Jeddah, Jeddah, Saudi Arabia

<sup>2</sup>Chemistry Department, Faculty of Applied Science, Taiz University, Taiz, Yemen

<sup>3</sup>Chemistry Department, Faculty of Science, University of Jeddah, Jeddah, Saudi Arabia

<sup>4</sup>Department of Biochemistry, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>5</sup>Center of Nanotechnology, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>6</sup>Department of biology, Faculty of Science, University of Jeddah, Jeddah, Saudi Arabia

<sup>7</sup>Stem Cell P2 Laboratory, The Center for Reproductive Medicine, Shantou University Medical College, Shantou, 515041, People's Republic of China

<sup>8</sup>Chemistry Department, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

<sup>9</sup>Biochemistry Department, Faculty of Veterinary Medicine, Zagazig University, Egypt

### Supplementary Table 4

Inhibitor effects on soluble and immobilized  $\alpha$ -amylase.

Inhibitor 2mM	Soluble $\alpha$ -amylase	OD at 560 nm n/3	Immobilized $\alpha$ -amylase	OD at 560 nm n/3
	Residual activity %		Residual activity %	
EDTA	18	0.219	60	0.438
Sodium Citrate	45	0.548	70	0.511
Sodium Oxalate	59	0.719	76	0.554
1,10 phenatroline	60	0.731	88	0.642
DTNB	61	0.745	84	0.613